

R-companion to: Estimation of the Thurstonian model for the 2-AC protocol

Rune Haubo Bojesen Christensen,
Hye-Seong Lee &
Per Bruun Brockhoff

May 16, 2011

This document describes how the examples in “Estimation of the Thurstonian model for the 2-AC protocol” (Christensen et al., 2011) can be executed in the free statistical software R (R Development Core Team, 2011) using the free R packages `sensR` and `ordinal` (Christensen, 2011; Christensen and Brockhoff, 2011) developed by the authors.

1 Example 1: Estimation of d' and τ

It is assumed that we have $\mathbf{n} = (2, 2, 6)$ observations in the three categories. We may estimate τ , d' and their standard errors with the `twoAC` function in the `sensR` package:

```
> library(sensR)
> fit <- twoAC(c(2, 2, 6))
> fit
```

Results for the 2-AC protocol with data `c(2, 2, 6)`:

	Estimate	Std. Error
tau	0.4159726	0.2674304
d.prime	0.7742595	0.5416717

Two-sided 95% confidence interval for d-prime based on the likelihood root statistic:

	Lower	Upper
d.prime	-0.2709832	1.859251

Significance test:

Likelihood root statistic = 1.446718 p-value = 0.14798
Alternative hypothesis: d-prime is different from 0

Alternatively we may compute τ and d' manually:

```
> n <- c(2, 2, 6)
> gamma <- cumsum(n/sum(n))
> z <- qnorm(gamma)[-3]
```

```

> z <- z * sqrt(2)
> (tau <- (z[2] - z[1])/2)

[1] 0.4159726

> (d <- -z[1] - tau)

[1] 0.7742595

```

2 Example 2: Inference for d-prime

The likelihood based confidence intervals and the one-sided discrimination significance test where the null hypothesis is “no sensory difference”, i.e., $d'_0 = 0$ using the likelihood root statistic are immediately available using the `twoAC` function:

```

> twoAC(c(2, 2, 6), d.prime0 = 0, conf.level = 0.95, statistic = "likelihood",
        alternative = "greater")

```

Results for the 2-AC protocol with data `c(2, 2, 6)`:

	Estimate	Std. Error
tau	0.4159726	0.2674304
d.prime	0.7742595	0.5416717

Two-sided 95% confidence interval for d-prime based on the likelihood root statistic:

	Lower	Upper
d.prime	-0.2709832	1.859251

Significance test:

Likelihood root statistic = 1.446718 p-value = 0.073988
 Alternative hypothesis: d-prime is greater than 0

The relative profile likelihood and Wald approximation can be obtained with:

```

> pr <- profile(fit)
> plot(pr)
> z <- pr$d.prime$d.prime
> w <- (coef(fit)[2, 1] - z)/coef(fit)[2, 2]
> lines(z, exp(-w^2/2), lty = 2)

```

3 Example 3: Power calculations

The function `twoACpwr` computes the power for the 2-AC protocol and a significance test of the users choice. The power assuming $\tau = 0.5$, $d' = 1$, the sample size is $N = 20$ for a two-sided preference test with $\alpha = 0.5$ is found with:

```

> twoACpwr(tau = 0.5, d.prime = 1, size = 20, d.prime0 = 0, alpha = 0.05,
           alternative = "two.sided", tol = 1e-05)

```

	power	actual.alpha	samples	discarded	kept	p.1	p.2	p.3
1	0.777677	0.04960103	231	94	137	0.1444	0.2174	0.6382




Figure 1: Relative profile likelihood curve (solid) and Wald approximation (dashed) for the data in example 1 and 2. The horizontal lines indicate 95% and 99% confidence intervals based on the likelihood function and the Wald approximation.

Apart from the power, we are told that the actual size of the test, α is close to the nominal 5%. The reason that the two differ is due to the discreteness of the observations, and hence the test statistic. We are also told that with $N = 20$ there are 231 possible outcomes of the 2-AC protocol. In computing the power 94 of these are discarded and power is computed based on the remaining 137 samples. The fraction of samples that are discarded is determined by the `tol` parameter. If we set this to zero, then no samples are discarded, however, the increase in precision is irrelevant:

```
> twoACpwr(tau = 0.5, d.prime = 1, size = 20, d.prime0 = 0, alpha = 0.05,
  alternative = "two.sided", tol = 0)

      power actual.alpha samples discarded kept    p.1    p.2    p.3
1 0.7776788  0.04960103      231          0  231 0.1444 0.2174 0.6382
```

The last three numbers in the output are the cell probabilities, so with $\tau = 0.5$ and $d' = 1$ we should, for example, expect around 22% of the answers in the “no difference” or “no preference” category.

4 Example 4: Estimation and standard errors via cumulative probit models

Estimates of τ and d' and their standard errors can be obtained from a cumulative probit model. We begin by defining the three leveled response variable `resp` and fitting a cumulative link model (CLM) using the function `clm` in package `ordinal` with weights equal to the observed counts and a probit link. Standard output contains the following coefficient table:

```

> library(ordinal)
> response <- gl(3, 1)
> fit.clm <- clm(response ~ 1, weights = c(2, 2, 6), link = "probit")
> (tab <- coef(summary(fit.clm)))

      Estimate Std. Error   z value   Pr(>|z|)
1|2 -0.8416213   0.4518154 -1.862755 0.06249676
2|3 -0.2533472   0.4009896 -0.631805 0.52751430

```

The τ and d' estimates are obtained by:

```

> theta <- tab[, 1]
> (tau <- (theta[2] - theta[1])/sqrt(2))

      2|3
0.4159726

> (d.prime <- (-theta[2] - theta[1])/sqrt(2))

      2|3
0.7742597

```

The variance-covariance matrix of the parameters can be extracted by the `vcov` method and the standard errors computed via the provided formulas:

```

> VCOV <- vcov(fit.clm)
> (se.tau <- sqrt((VCOV[1, 1] + VCOV[2, 2] - 2 * VCOV[2, 1])/2))

[1] 0.2674311

> (se.d.prime <- sqrt((VCOV[1, 1] + VCOV[2, 2] + 2 * VCOV[2, 1])/2))

[1] 0.5416737

```

Observe how these estimates and standard errors are identical to those in Example 1.

We could also have used the `clm2twoAC` function from package `sensR` which extract estimates and standard errors from a `clm` model fit object:

```

> clm2twoAC(fit.clm)

      Estimate Std. Error   z value   Pr(>|z|)
tau      0.4159726   0.2674311  1.555439  0.11984
d-prime  0.7742597   0.5416737  1.429384  0.15289

```

5 Example 5: A regression model for d'

Assume that a study is performed and gender differences in the discrimination ability is of interest. Suppose that (20, 20, 60) is observed for women and (10, 20, 70) is observed for men. The standard output from a cumulative probit model contains the coefficient table:

```

> n.women <- c(2, 2, 6) * 10
> n.men <- c(1, 2, 7) * 10
> wt <- c(n.women, n.men)
> response <- gl(3, 1, length = 6)
> gender <- gl(2, 3, labels = c("women", "men"))

```

```
> fm2 <- clm(response ~ gender, weights = wt, link = "probit")
> (tab2 <- coef(summary(fm2)))
```

	Estimate	Std. Error	z value	Pr(> z)
1 2	-0.8887474	0.1346051	-6.602627	4.039349e-11
2 3	-0.2283095	0.1238663	-1.843194	6.530069e-02
gendermen	0.3205403	0.1741445	1.840658	6.567176e-02

The estimate of τ (assumed constant between genders) and the gender specific estimates of d' can be extracted by:

```
> theta <- fm2$Theta
> (tau <- (theta[2] - theta[1])/sqrt(2))

2|3
0.4670001

> (d.women <- (-theta[2] - theta[1])/sqrt(2))

2|3
0.7898785

> (d.men <- d.women + fm2$beta * sqrt(2))

2|3
1.243191
```

Again we could use the `clm2twoAC` function to get the coefficient table for the 2-AC model from the CLM-model:

```
> clm2twoAC(fm2)

Estimate Std. Error z value Pr(>|z|)
tau      0.4670001 0.06700014 6.970135 3.1664e-12
d-prime  0.7898785 0.17021271 4.640538 3.4750e-06
gendermen 0.4533125 0.24627746 1.840658 0.065672
```

Observe that d' for women is given along with the difference in d' for men and women rather than d' for each of the genders.

The Wald test for gender differences is directly available from the coefficient table with $p = 0.0657$. The corresponding likelihood ratio test can be obtained by:

```
> anova(fm2, update(fm2, location = ~. - gender))

Likelihood ratio tests of cumulative link models

Response: response
      Model Resid. df -2logLik  Test   Df LR stat.   Pr(Chi)
1      1 | |      198 354.5858
2 gender | |      197 351.1861 1 vs 2    1   3.39971 0.0652079
```

which is slightly closer to significance. The 95% profile likelihood confidence interval for the difference between men and women on the d' -scale is:

```
> confint(fm2) * sqrt(2)

2.5 %    97.5 %
loc.gendermen -0.02850948 0.9372278
```

6 Regression model for replicated 2-AC data

The data used in example 6 are not directly available at the time of writing. Assume however, that data are available in the R data.frame `repData`. Then the cumulative probit mixed model where preference depends on reference and consumers (`cons`) are random can be fitted with:

```
> fm3.agq <- clmm(preference ~ reference, random = cons, nAGQ = 10,
  data = repData, link = "probit", Hess = TRUE)
> summary(fm3.agq)
```

Cumulative Link Mixed Model fitted with the adaptive Gauss-Hermite quadrature approximation with 10 quadrature points

Call:

```
clmm(location = preference ~ reference, random = cons, data = repData,
  Hess = TRUE, link = "probit", nAGQ = 10)
```

Random effects:

	Var	Std.Dev
cons	1.367068	1.169217

Location coefficients:

	Estimate	Std. Error	z value	Pr(> z)
referenceB	0.4059	0.1002	4.0519	5.0806e-05

No scale coefficients

Threshold coefficients:

	Estimate	Std. Error	z value
A N	0.4823	0.1130	4.2680
N B	0.8488	0.1166	7.2801

log-likelihood: -668.9122

AIC: 1345.824

Condition number of Hessian: 36.79649

Here we asked for the accurate 10-point adaptive Gauss-Hermite quadrature approximation. The 2-AC estimates are available using the `clm2twoAC` function:

```
> clm2twoAC(fm3.agq)
```

	Estimate	Std. Error	z value	Pr(> z)
tau	0.2592190	0.02902614	8.930538	< 2.22e-16
d-prime	-0.9412295	0.15975045	-5.891874	3.8184e-09
referenceB	0.5740292	0.14166951	4.051889	5.0806e-05

The standard deviation of the consumer-specific d 's is given by:

```
> fm3.agq$stDev * sqrt(2)
```

	Std.Dev
cons	1.653522

The profile likelihood curve can be obtained using the profile method:

```
> pr <- profile(fm3.agq, range = c(0.7, 1.8))
```




Figure 2: Profile likelihood for σ_d . Horizontal lines indicate 95% and 99% confidence limits.

And then plottet using:

```
> plpr <- plot(pr, fig = FALSE)
> plot(sqrt(2) * plpr$stDev$x, plpr$stDev$y, type = "l", xlab = expression(sigma[d]),
       ylab = "Relative profile likelihood", xlim = c(1, 2.5), axes = FALSE)
> axis(1)
> axis(2, las = 1)
> abline(h = attr(plpr, "limits"))
> text(2.4, 0.17, "95% limit")
> text(2.4, 0.06, "99% limit")
```

The resulting figure is shown in Figure 2. The profile likelihood confidence intervals are obtained using:

```
> confint(pr, level = 0.95) * sqrt(2)

      2.5 %    97.5 %
stDev 1.362329 2.001483
```

The probabilities that consumers prefer yoghurt A, have no preference or prefer yoghurt B can, for an average consumer be obtained by predicting from the CLMM:

```
> newdat <- expand.grid(preference = factor(c("A", "N", "B"), levels = c("A",
  "N", "B"), ordered = TRUE), reference = factor(c("A", "B")))
> pred <- predict(fm3.agq, newdata = newdat)
```

The predictions for the extreme consumers have to be obtained by hand. Here we ask for the predictions for the 5th percentile and 95th percentile of the consumer population for reference A and B:

```
> q95.refA <- diff(c(0, pnorm(fm3.agq$Theta - qnorm(1 - 0.05) *
  fm3.agq$stDev), 1))
> q05.refA <- diff(c(0, pnorm(fm3.agq$Theta - qnorm(0.05) * fm3.agq$stDev),
  1))
```

```

> q95.refB <- diff(c(0, pnorm(fm3.agq$Theta - fm3.agq$beta - qnorm(1 -
  0.05) * fm3.agq$stDev), 1))
> q05.refB <- diff(c(0, pnorm(fm3.agq$Theta - fm3.agq$beta - qnorm(0.05) *
  fm3.agq$stDev), 1))

```

Plotting follows the standard methods:

```

> par(mar = c(0, 2, 0, 0.5) + 0.5)
> plot(1:3, pred[1:3], ylim = c(0, 1), axes = FALSE, xlab = "",
  ylab = "", pch = 19)
> axis(1, lwd.ticks = 0, at = c(1, 3), labels = c("", ""))
> axis(2, las = 1)
> points(1:3, pred[4:6], pch = 1)
> lines(1:3, pred[1:3])
> lines(1:3, pred[4:6], lty = 2)
> text(2, 0.6, "Average consumer")
> legend("topright", c("Reference A", "Reference B"), lty = 1:2,
  pch = c(19, 1), bty = "n")

> par(mar = c(0, 2, 0, 0.5) + 0.5)
> plot(1:3, q05.refA, ylim = c(0, 1), axes = FALSE, xlab = "",
  ylab = "", pch = 19)
> axis(1, lwd.ticks = 0, at = c(1, 3), labels = c("", ""))
> axis(2, las = 1)
> points(1:3, q05.refB, pch = 1)
> lines(1:3, q05.refA)
> lines(1:3, q05.refB, lty = 2)
> text(2, 0.6, "5th percentile consumer")

> par(mar = c(2, 2, 0, 0.5) + 0.5)
> plot(1:3, q95.refA, ylim = c(0, 1), axes = FALSE, xlab = "",
  ylab = "", pch = 19)
> axis(1, at = 1:3, labels = c("prefer A", "no preference", "prefer B"))
> axis(2, las = 1)
> points(1:3, q95.refB, pch = 1)
> lines(1:3, q95.refA)
> lines(1:3, q95.refB, lty = 2)
> text(2, 0.6, "95th percentile consumer")

```

The resulting figure is shown in Fig. 3.

7 End note

Version details for R and the packages `sensR` and `ordinal` appear below:

```

> sessionInfo()

R version 2.13.0 (2011-04-13)
Platform: i386-pc-mingw32/i386 (32-bit)

locale:
[1] LC_COLLATE=Danish_Denmark.1252 LC_CTYPE=Danish_Denmark.1252
[3] LC_MONETARY=Danish_Denmark.1252 LC_NUMERIC=C
[5] LC_TIME=Danish_Denmark.1252

```





Figure 3: The probabilities of preferring yoghurt A, having no preference, or preferring yoghurt B for an average consumer ($b = 0$) and for fairly extreme consumers ($b = \pm 1.64\sigma_b$).

attached base packages:

```
[1] stats      graphics  grDevices  utils      datasets  methods   base
```

other attached packages:

```
[1] sensR_1.2.8      ordinal_2011.05-10 ucminf_1.1-0      numDeriv_2010.11-1
[5] MASS_7.3-13
```

References

- Christensen, R. H. B. (2011). ordinal—regression models for ordinal data. R package version 2010.10-22 <http://www.cran.r-project.org/package=ordinal/>.
- Christensen, R. H. B. and P. B. Brockhoff (2011). sensR: An R-package for Thurstonian modelling of discrete sensory data. R package version 1.2.8 <http://www.cran.r-project.org/package=sensR/>.
- Christensen, R. H. B., H.-S. Lee, and P. B. Brockhoff (2011). Estimation of the Thurstonian model for the 2-AC protocol. *Submitted to Food Quality and Preference*.
- R Development Core Team (2011). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. ISBN 3-900051-07-0.