

Package ‘LIStest’

October 12, 2022

Type Package

Title Tests of independence based on the Longest Increasing Subsequence

Version 2.1

Date 2014-03-12

Author J. E. Garcia and V. A. Gonzalez-Lopez

Maintainer J. E. Garcia <jg@ime.unicamp.br>

Depends R (>= 2.10)

Description Tests for independence between X and Y computed from a paired sample $(x_1, y_1), \dots, (x_n, y_n)$ of (X, Y) , using one of the following statistics (a) the Longest Increasing Subsequence (L_n), (b) JL_n , a Jackknife version of L_n or (c) JLM_n , a Jackknife version of the longest monotonic subsequence. This family of tests can be applied under the assumption of continuity of X and Y.

License GPL-2

LazyLoad yes

LazyData yes

NeedsCompilation no

Repository CRAN

Date/Publication 2014-03-12 23:17:20

R topics documented:

LIStest-package	2
JLMn	2
JLn	3
lis	5
lis.test	6
Ln	7
TJLMN	8
TJLN	9
TLN	9

Index

10

<code>LITest-package</code>	<i>Tests of independence based on the Longest Increasing Subsequence</i>
-----------------------------	--

Description

Tests for independence between X and Y computed from a paired sample $(x_1, y_1), \dots, (x_n, y_n)$ of (X, Y) , using one of the following statistics (a) the Longest Increasing Subsequence (L_n), (b) JL_n , a Jackknife version of L_n or (c) JLM_n , a Jackknife version of the longest monotonic subsequence. This family of tests can be applied under the assumption of continuity of X and Y.

Details

Package:	LIStest
Type:	Package
Version:	2.1
Date:	2014-03-12
License:	GPL-2

Author(s)

J. E. Garcia and V. A. Gonzalez-Lopez Maintainer: J. E. Garcia <jg@ime.unicamp.br>

References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

<code>JLMn</code>	<i>JLMn statistic, to test independence</i>
-------------------	---

Description

It compute the JLM_n -statistic, from a bivariate sample of continuous random variables X and Y.

Usage

`JLMn(x, y)`

Arguments

<code>x, y</code>	numeric vectors of data values. x and y must have the same length.
-------------------	--

Details

See subsection 3.3-Main reference. For sample sizes less than 20, the correction introduced in subsection 3.2 from main reference, with $c = 0.4$ was avoided.

Value

The value of the JLMn-statistic.

Author(s)

J. E. Garcia, V. A. Gonzalez-Lopez

References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

Examples

```
# mixture of two bivariate normal, one with correlation 0.9 and
# the other with correlation -0.9
#
N <-100
ro<- 0.90
Z1<-rnorm(N)
Z2<-rnorm(N)
X2<-X1<-Z1
I<-(1:floor(N*0.5))
I2<-(floor(N*0.5)+1):N
X1[I]<-Z1[I]
X2[I]<-(Z1[I]*ro+Z2[I]*sqrt(1-ro*ro))
X1[I2]<-Z1[I2]
X2[I2]<-(Z1[I2]*(-ro)+Z2[I2]*sqrt(1-ro*ro))
plot(X1,X2)

#calculate the statistic
a<-JLMn(X1,X2)
a
```

Description

It compute the JL_n-statistic, from a bivariate sample of continuous random variables X and Y.

Usage

```
JLn(x, y)
```

Arguments

x, y	numeric vectors of data values. x and y must have the same length.
------	--

Details

See subsection 3.2.-Main reference. For sample sizes less than 20, the correction introduced in subsection 3.2 from main reference, with $c = 0.4$ was avoided.

Value

The value of the JLn-statistic.

Author(s)

J. E. Garcia and V. A. Gonzalez-Lopez

References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

Examples

```
## mixture of two bivariate normal, one with correlation 0.9 and
## the other with correlation -0.9
#
N <- 100
ro<- 0.90
Z1<-rnorm(N)
Z2<-rnorm(N)
X2<-X1<-Z1
I<-(1:floor(N*0.5))
I2<-(floor(N*0.5)+1):N
X1[I]<-Z1[I]
X2[I]<-(Z1[I]*ro+Z2[I]*sqrt(1-ro*ro))
X1[I2]<-Z1[I2]
X2[I2]<-(Z1[I2]*(-ro)+Z2[I2]*sqrt(1-ro*ro))
plot(X1,X2)

# calculate the statistic
a<-JLn(X1,X2)
a
```

lis*Longest increasing subsequence for a univariate sample*

Description

It compute the size of the longest increasing subsequence from a sample of a (continuous) random variable.

Usage

```
lis(x)
```

Arguments

x numeric vector of data values.

Details

See example 2.1-Main reference.

Value

Integer, the size of the longest increasing subsequence.

Author(s)

J. E. Garcia and V. A. Gonzalez-Lopez

References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

Examples

```
#see Example 2.1 (reference)
a<-lis(c(3,6,1,7,4,2,5,8))
a
```

lis.test*Test for independence between paired samples*

Description

Test for independence between X and Y computed from a paired sample ($x_1, y_1, \dots, x_n, y_n$) of (X, Y), using one of the following statistics (a) the Longest Increasing Subsequence (Ln), (b) JLn, a Jackknife version of Ln or (c) JLMn, a Jackknife version of the longest monotonic subsequence. This family of tests can be applied under the assumption of continuity of X and Y.

Usage

```
lis.test(x, y, alternative = c("two.sided", "less", "greater"),
method = c("JLMn", "Ln", "JLn"))
```

Arguments

- | | |
|--------------------------|--|
| <code>x, y</code> | numeric vectors of data values. x and y must have the same length. |
| <code>alternative</code> | indicates the alternative hypothesis and must be one of "two.sided"(default), "greater" or "less". |
| <code>method</code> | a character string indicating which statistics is to be used for the test. One of "Ln", "JLn", or "JLMn"(default). |

Details

For sample sizes less than 20, the correction introduced in subsection 3.2 from main reference, with $c = 0.4$ was avoided.

Value

- | | |
|------------------------------|--|
| <code>sample.estimate</code> | the value of the statistic. |
| <code>p.value</code> | the p-value for the test. |
| <code>alternative</code> | a character string describing the alternative hypothesis. |
| <code>method</code> | a character string indicating what type of Lis-test was performed. |

Author(s)

J. E. Garcia and V. A. Gonzalez-Lopez

References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

Examples

```

# Example 1
# mixture of two bivariate normal, one with correlation 0.9
# and the other with correlation -0.9

N <-100
ro<- 0.90
Z1<-rnorm(N)
Z2<-rnorm(N)
X2<-X1<-Z1
I<-(1:floor(N*0.5))
I2<-(floor(N*0.5)+1):N
X1[I]<-Z1[I]
X2[I]<-(Z1[I]*ro+Z2[I]*sqrt(1-ro*ro))
X1[I2]<-Z1[I2]
X2[I2]<-(Z1[I2]*(-ro)+Z2[I2]*sqrt(1-ro*ro))
plot(X1,X2)
# calculate the p.value using the default settings (method="JLMn"
# and alternative="two.sided")
lis.test(X1,X2)
# calculate the p.value using method="JLn" and
# alternative="two.sided".
lis.test(X1,X2,method="JLn")
#
# Example 2: see subsection 4.3.2-Application 2 from main reference.
# (It requires the package VGAM)
#
#require(VGAM)
#plot(coalminers$BW, coalminers$nBW)
#lis.test(coalminers$BW, coalminers$nBW,
#alternative = "greater", method = "Ln")
#lis.test(coalminers$BW, coalminers$nBW,
#alternative = "greater", method = "JLn")
#

```

Ln

Ln (Longest Increasing Subsequence) statistic, to test independence

Description

It compute the Ln-statistic, from a bivariate sample of continuous random variables X and Y.

Usage

`Ln(x, y)`

Arguments

<code>x, y</code>	numeric vectors of data values. x and y must have the same length.
-------------------	--

Details

See Section 2.-Main reference.

Value

The value of the Ln-statistic.

Author(s)

J. E. Garcia and V. A. Gonzalez-Lopez

References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

Examples

```
## mixture of two bivariate normal, one with correlation
## 0.9 and the other with correlation -0.9
#
N <-100
ro<- 0.90
Z1<-rnorm(N)
Z2<-rnorm(N)
X2<-X1<-Z1
I<-(1:floor(N*0.5))
I2<-(floor(N*0.5)+1):N
X1[I]<-Z1[I]
X2[I]<-(Z1[I]*ro+Z2[I]*sqrt(1-ro*ro))
X1[I2]<-Z1[I2]
X2[I2]<-(Z1[I2]*(-ro)+Z2[I2]*sqrt(1-ro*ro))
plot(X1,X2)

# calculate the statistic
a<-Ln(X1,X2)
a
```

Description

Simulated values for the JLMn statistic under the hypothesis of independence

Format

The format is: List of 200 tables

References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

TJLN	<i>Simulated values for the JLn statistic</i>
------	---

Description

Simulated values for the JLn statistic under the hypothesis of independence.

Format

The format is: List of 200 tables

References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

TLN	<i>Simulated values for the Ln statistic</i>
-----	--

Description

Simulated values for the Ln statistic under the hypothesis of independence

Format

The format is: List of 200 tables

References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

Index

* ~copula

JLMn, 2

JLn, 3

lis, 5

lis.test, 6

Ln, 7

* ~longest increasing subsequence

JLMn, 2

JLn, 3

lis, 5

lis.test, 6

Ln, 7

* datasets

TJLMN, 8

TJLN, 9

TLN, 9

JLMn, 2

JLn, 3

lis, 5

lis.test, 6

LITest (LITest-package), 2

LITest-package, 2

Ln, 7

TJLMN, 8

TJLN, 9

TLN, 9