

Package ‘ecotourism’

September 16, 2025

Type Package

Title Collection of Data on Wildlife Sightings, Tourism Counts, and Weather from Australia

Version 0.1.0

Description This is a collection of data files for exploring sightings of wild things, relative to weather and tourism patterns in Australia.

License MIT + file LICENSE

Encoding UTF-8

LazyData true

LazyDataCompression xz

Suggests dplyr, ggplot2, ggthemes, ggbeeswarm, sf, knitr, lubridate, quarto, testthat (>= 3.0.0), tidyr

VignetteBuilder quarto

SystemRequirements Quarto

Depends R (>= 3.5)

URL <https://github.com/vahdatjavad/ecotourism>

BugReports <https://github.com/vahdatjavad/ecotourism/issues>

RoxygenNote 7.3.2

Config/testthat/edition 3

NeedsCompilation no

Author Dianne Cook [aut] (ORCID: <<https://orcid.org/0000-0002-3813-7155>>),
Lyn Cook [aut] (ORCID: <<https://orcid.org/0000-0002-3172-4920>>),
Javad Vahdat Atashgah [aut, cre] (ORCID:
<<https://orcid.org/0009-0004-5845-2584>>)

Maintainer Javad Vahdat Atashgah <vahdatjavad@gmail.com>

Repository CRAN

Date/Publication 2025-09-16 06:00:11 UTC

Contents

glowworms	2
gouldian_finch	3
manta_rays	4
orchids	5
oz_lga	6
top_stations	7
tourism_quarterly	7
tourism_region	8
weather	9
weather_stations	10
Index	12

glowworms	<i>Glowworms Occurrence Data (2014–2024)</i>
-----------	--

Description

This dataset contains cleaned and enriched occurrence records for glowworms (*Arachnocampa tasmaniensis*) in Australia from 2014 to 2024. It includes spatial, temporal, taxonomic, and weather station metadata.

Usage

glowworms

Format

- A tibble with 124 rows and 14 variables:
- obs_lat** Latitude of the observation (decimal degrees)
 - obs_lon** Longitude of the observation (decimal degrees)
 - date** Observation date (YYYY-MM-DD)
 - time** Observation time (HH:MM:SS, character)
 - year** Observation year
 - month** Month of the observation
 - day** Day of the month
 - hour** Hour of the day (0–23)
 - weekday** Day of the week (ordered factor)
 - dayofyear** Day of the year (1–366)
 - sci_name** Scientific name of the observed species
 - record_type** Type of observation (e.g., HUMAN_OBSERVATION)
 - obs_state** Australian state where the observation occurred
 - ws_id** ID of the nearest weather station (e.g., "949610-99999")

Details

Data was sourced via the ‘galah’ package from the Atlas of Living Australia, filtered and cleaned, then enriched by linking each record to the nearest weather station using geospatial methods.

Source

Atlas of Living Australia via **galah**

Examples

```
data(glowworms)
head(glowworms)
```

gouldian_finch	<i>Gouldian Finch Occurrence Data (2014–2024)</i>
----------------	---

Description

This dataset contains cleaned and processed occurrence records for the Gouldian Finch (*Chloebeia gouldiae*) in Australia between 2014 and 2024. It includes spatial coordinates, temporal details, species information, and the ID of the nearest weather station for each observation.

Usage

```
gouldian_finch
```

Format

- A tibble with 3,921 rows and 14 variables:
- obs_lat** Latitude of the observation (decimal degrees)
 - obs_lon** Longitude of the observation (decimal degrees)
 - date** Date of the observation (YYYY-MM-DD)
 - time** Time of the observation (HH:MM:SS)
 - year** Year of the observation
 - month** Month (1–12)
 - day** Day of the month
 - hour** Hour extracted from the time (0–23)
 - weekday** Day of the week (as ordered factor)
 - dayofyear** Day of the year (1–366)
 - sci_name** Scientific name of the species
 - record_type** Type of observation (e.g., HUMAN_OBSERVATION)
 - obs_state** Australian state where the observation was recorded
 - ws_id** Nearest weather station ID (e.g., "948280-99999")

Details

The data was retrieved from the Atlas of Living Australia using the **galah** package, then standardized, cleaned, and matched to the three closest weather stations using geospatial tools.

Source

Atlas of Living Australia via **galah**

See Also

[glowworms](#), [weather](#)

Examples

```
data(gouldian_finch)
head(gouldian_finch)
```

manta_rays	<i>Manta Ray Occurrence Data (2014–2024)</i>
------------	--

Description

This dataset contains occurrence records for the reef manta ray (**Mobula alfredi**) observed in Australian waters from 2014 to 2024. The data includes spatial and temporal metadata, species identifiers, and linked weather station IDs.

Usage

```
manta_rays
```

Format

- A tibble with 1,088 rows and 14 variables:
- obs_lat** Latitude of the observation (decimal degrees)
 - obs_lon** Longitude of the observation (decimal degrees)
 - date** Date of the observation (YYYY-MM-DD)
 - time** Time of the observation (HH:MM:SS)
 - year** Year of the observation
 - month** Month (1–12)
 - day** Day of the month
 - hour** Hour extracted from the time (0–23)
 - weekday** Day of the week (as ordered factor)
 - dayofyear** Day of the year (1–366)
 - sci_name** Scientific name — all observations are *Mobula alfredi*

- record_type** Type of observation (e.g., MACHINE_OBSERVATION)
- obs_state** Australian state where the observation occurred (may be missing)
- ws_id** Nearest weather station ID (e.g., "947800-99999")

Details

Records were accessed using the **galah** package and filtered specifically for **Mobula alfredi**. Data has been cleaned and enriched with spatial proximity to weather stations for climate-related analysis.

Source

Atlas of Living Australia via **galah**

See Also

[gouldian_finch](#), [weather](#)

Examples

```
data(manta_rays)
head(manta_rays)
```

orchids	<i>Orchid Occurrence Data (2014–2024)</i>
---------	---

Description

This dataset contains over 300,000 occurrence records of orchid species (**Orchidaceae**) in Australia from 2014 to 2024. The data includes spatial, temporal, and taxonomic details, as well as associated weather station metadata for ecological analysis.

Usage

```
orchids
```

Format

- A tibble with 302,123 rows and 14 variables:
- obs_lat** Latitude of the observation (decimal degrees)
 - obs_lon** Longitude of the observation (decimal degrees)
 - date** Date of the observation (YYYY-MM-DD)
 - time** Time of the observation (HH:MM:SS)
 - year** Year of the observation
 - month** Month (1–12)

day Day of the month
hour Hour extracted from the time (0–23)
weekday Day of the week (as ordered factor)
dayofyear Day of the year (1–366)
sci_name Scientific name of the observed orchid species
record_type Type of observation (e.g., HUMAN_OBSERVATION, PRESERVED_SPECIMEN)
obs_state Australian state where the observation occurred (may be missing)
ws_id Nearest weather station ID linked to the observation

Details

The data was collected using the **galah** package from the Atlas of Living Australia, cleaned, and linked to nearby weather stations for ecological and climatic studies. The records span multiple orchid genera and include a range of observation types.

Source

Atlas of Living Australia via **galah**

See Also

[glowworms](#), [gouldian_finch](#), [weather](#)

Examples

```
data(orchids)
head(orchids)
```

oz_lga	<i>oz_lga</i>
--------	---------------

Description

LGA polygons for Australia

Usage

```
oz_lga
```

Format

A spatial polygon object

Examples

```
head(oz_lga)
```

top_stations	<i>Top Weather Stations for Each Organism</i>
--------------	---

Description

A lookup table identifying the top 3 most frequently linked weather stations for each focal organism in the ecotourism package. These stations were selected based on the number of linked observations across a 10-year period (2014–2024).

Usage

top_stations

Format

A tibble with 12 rows and 2 variables:

- organism** Name of the organism (e.g., "glowworms", "orchids")
- ws_id** Weather station ID (e.g., "948720-99999")

Details

This table was created by counting the frequency of ‘ws_id’ assignments within each organism dataset and selecting the top 3 stations per organism. These top stations are used for downloading daily weather data via the **GSODR** package.

See Also

[weather](#), [weather_stations](#)

Examples

```
data(top_stations)
head(top_stations)
```

tourism_quarterly	<i>Quarterly Tourism Trips by Region and Purpose</i>
-------------------	--

Description

A dataset containing quarterly estimates of overnight tourism trips in Australia, broken down by trip purpose and tourism region.

Usage

tourism_quarterly

Format

A data frame with `nrow(tourism_quarterly)` rows and 4 variables: `year`: The year of the tourism data `quarter`: Quarter number like 1, 2, 3, 4 `purpose`: Purpose of visit category: - "Holiday" - "Business" `trips`: Number of overnight trips (in thousands). `region_id`: Unique integer identifier linking to the `tourism_region` dataset. `ws_id`: Identifier of the nearest Bureau of Meteorology weather station to the tourism region.

Details

Tourism regions are formed through the aggregation of Statistical Local Areas (SLAs) or similar ABS-defined geographies, as determined by state and territory tourism authorities. This dataset is designed for analysis of seasonal tourism patterns and can be joined to `tourism_region` for spatial analysis.

References

Tourism Research Australia: <https://www.tra.gov.au>

Examples

```
data(tourism_quarterly)
head(tourism_quarterly)
```

tourism_region	<i>Tourism Regions and Nearest Weather Stations</i>
----------------	---

Description

A dataset containing the locations of Australian tourism regions, their geographic coordinates, and the nearest Bureau of Meteorology weather station. Each region is assigned a unique identifier for linking to other tourism datasets.

Usage

```
tourism_region
```

Format

A data frame with `nrow(tourism_region)` rows and 5 variables: `region`: Name of the tourism region. Tourism regions are defined by Tourism Research Australia and generally formed through the aggregation of Statistical Local Areas (SLAs) or other ABS-defined geographies. `lon`: Longitude of the tourism region representative point (WGS84). `lat`: Latitude of the tourism region representative point (WGS84). `region_id`: Unique integer identifier for the tourism region. Useful for joining with other tourism-related datasets. `ws_id`: Identifier of the nearest Bureau of Meteorology weather station to the tourism region.

Details

Coordinates for each tourism region are intended to represent a central location within the region (e.g., polygon centroid). The nearest weather station is determined using great-circle distance calculations based on the Bureau of Meteorology’s official station list.

References

Tourism Research Australia: <https://www.tra.gov.au> Australian Bureau of Meteorology: <http://www.bom.gov.au>

Examples

```
data(tourism_region)
head(tourism_region)
```

weather	<i>Daily Weather Data for Top Stations (2014–2024)</i>
---------	--

Description

This dataset contains daily weather observations for the top weather stations associated with focal species in the ecotourism package. Data spans from 2014 to 2024 and includes temperature, humidity, precipitation, and wind measures.

Usage

```
weather
```

Format

- A tibble with 35,527 rows and 18 variables:
- ws_id** Weather station ID (e.g., "948720-99999")
 - stn_lat** Latitude of the weather station
 - stn_lon** Longitude of the weather station
 - date** Observation date (YYYY-MM-DD)
 - year** Year of observation
 - month** Month of observation (1–12)
 - day** Day of the month
 - weekday** Day of the week (as ordered factor)
 - dayofyear** Day of the year (1–366)
 - temp** Average temperature (°C)
 - min** Minimum temperature (°C)
 - max** Maximum temperature (°C)

- dewp** Dew point temperature (°C)
- rh** Relative humidity (%)
- prcp** Precipitation (mm)
- rainy** Binary flag indicating whether PRCP > 5 mm (1 = rainy day)
- wind_speed** Average wind speed (m/s)
- max_speed** Maximum sustained wind speed (m/s)

Details

The weather data was retrieved from the Global Surface Summary of the Day (GSOD) dataset via the **GSODR** package for the top 3 weather stations per organism in the ecotourism project. This data supports climate-biodiversity interaction analyses.

Source

GSOD via **GSODR**

See Also

[top_stations](#), [glowworms](#), [gouldian_finch](#), [weather_stations](#)

Examples

```
data(weather)
head(weather)
```

weather_stations	<i>Australian Weather Station Metadata</i>
------------------	--

Description

This dataset contains metadata for 732 weather stations across Australia, including coordinates, station names, and geocoded location details.

Usage

```
weather_stations
```

Format

- A tibble with 732 rows and 7 variables:
- ws_id** Weather station ID (e.g., "941000-99999")
 - stname** Station name (e.g., "KALUMBURU")
 - stn_lat** Latitude of the station (decimal degrees)
 - stn_lon** Longitude of the station (decimal degrees)
 - address** Full geocoded address (from reverse geocoding)
 - stn_city** Parsed city or locality name
 - stn_state** Australian state or territory

Details

This data was derived from the GSOD inventory using the **GSODR** package, filtered for Australian stations, and geocoded using OpenStreetMap APIs. It is used to match ecological observations with relevant local weather conditions.

Source

GSOD inventory via **GSODR**; geocoded with OpenStreetMap

See Also

[weather](#), [top_stations](#), [gouldian_finch](#)

Examples

```
data(weather_stations)
head(weather_stations)
```

Index

* datasets

- glowworms, [2](#)
- gouldian_finch, [3](#)
- manta_rays, [4](#)
- orchids, [5](#)
- oz_lga, [6](#)
- top_stations, [7](#)
- tourism_quarterly, [7](#)
- tourism_region, [8](#)
- weather, [9](#)
- weather_stations, [10](#)

glowworms, [2](#), [4](#), [6](#), [10](#)

gouldian_finch, [3](#), [5](#), [6](#), [10](#), [11](#)

manta_rays, [4](#)

orchids, [5](#)

oz_lga, [6](#)

top_stations, [7](#), [10](#), [11](#)

tourism_quarterly, [7](#)

tourism_region, [8](#), [8](#)

weather, [4–7](#), [9](#), [11](#)

weather_stations, [7](#), [10](#), [10](#)